

ON THE STRUCTURE OF COMPLEX HOMOGENEOUS SUPERMANIFOLDS

E.G. Vishnyakova ¹

Tver State University

ABSTRACT. For a Lie group G and a closed Lie subgroup $H \subset G$, it is well known that the coset space G/H can be equipped with the structure of a manifold homogeneous under G and that any G -homogeneous manifold is isomorphic to one of this kind. An interesting problem is to find an analogue of this result in the case of supermanifolds.

In the classical setting, G is a real or a complex Lie group and G/H is a real and, respectively, a complex manifold. Now, if G is a real Lie supergroup and $H \subset G$ is a closed Lie subsupergroup, there is a natural way to consider G/H as a supermanifold. Furthermore, any G -homogeneous real supermanifold can be obtained in this way, see [1]. The goal of this paper is to give a proof of this result in the complex case.

1. Preliminaries

We will use the word "supermanifold" in the sense of Beresin-Leites-Kostant (see [1, 2]). All the time, we will be interested in the complex-analytic version of the theory. A morphism $(M, \mathcal{O}_M) \rightarrow (N, \mathcal{O}_N)$ between two supermanifolds is denoted by $\varphi = (\varphi_1, \varphi_2)$, where $\varphi_1 : M \rightarrow N$ is a continuous mapping and $\varphi_2 : \mathcal{O}_N \rightarrow (\varphi_1)_*(\mathcal{O}_M)$ is a homomorphism of sheaves of superalgebras. We begin with the more general notion of a Lie supergroup.

A *Lie supergroup* is a supermanifold (G, \mathcal{O}_G) , for which the following three morphisms are defined: $\nu : (G, \mathcal{O}_G) \times (G, \mathcal{O}_G) \rightarrow (G, \mathcal{O}_G)$ (multiplication morphism), $\iota : (G, \mathcal{O}_G) \rightarrow (G, \mathcal{O}_G)$ (passing to the inverse), $\varepsilon : (\text{pt}, \mathbb{C}) \rightarrow (G, \mathcal{O}_G)$ (identity morphism). Moreover, these morphisms should satisfy the usual conditions, modeling the group axioms. The underlying manifold G is a Lie group. We denote by \mathfrak{g} the Lie superalgebra of (G, \mathcal{O}_G) (see [2] for the corresponding definition).

An *action of a Lie supergroup* (G, \mathcal{O}_G) on a supermanifold (M, \mathcal{O}_M) is a morphism $\mu : (G, \mathcal{O}_G) \times (M, \mathcal{O}_M) \rightarrow (M, \mathcal{O}_M)$, such that the following conditions hold:

- $\mu \circ (\nu \times \text{id}) = \mu \circ (\text{id} \times \mu)$;
- $\mu \circ (\varepsilon \times \text{id}) = \text{id}$.

¹Mathematics Classification Number 32C11

Key words and phrases. Complex homogeneous supermanifold, Lie supergroup, action of a Lie supergroup.

Work supported by German Academic Exchange Service, Michail-Lomonosov-Forschungstipendien and by the Russian Foundation for Basic Research (grant no. 07-01-00230).

We denote by $\mathfrak{v}(M, \mathcal{O}_M)$ the Lie superalgebra of holomorphic vector fields on (M, \mathcal{O}_M) . Let \mathfrak{m}_x be the maximal ideal of the local superalgebra $(\mathcal{O}_M)_x$. The vector superspace $T_x(M, \mathcal{O}_M) = (\mathfrak{m}_x/\mathfrak{m}_x^2)^*$ is called the *tangent space* to (M, \mathcal{O}_M) at $x \in M$. From the inclusions $v(\mathfrak{m}_x) \subset (\mathcal{O}_M)_x$ and $v(\mathfrak{m}_x^2) \subset \mathfrak{m}_x$, where $v \in \mathfrak{v}(M, \mathcal{O}_M)$, it follows that there exists an even linear mapping $\text{ev}_x(v) : \mathfrak{m}_x/\mathfrak{m}_x^2 \rightarrow (\mathcal{O}_M)_x/\mathfrak{m}_x \simeq \mathbb{C}$. In other words, $\text{ev}_x(v) \in T_x(M, \mathcal{O}_M)$, and so we obtain a map $\text{ev}_x : \mathfrak{v}(M, \mathcal{O}_M) \rightarrow T_x(M, \mathcal{O}_M)$.

Let $(U, \mathcal{O}_M) \subset (M, \mathcal{O}_M)$ be a superdomain with even and, respectively, odd coordinates (x_i) and (ξ_j) and let $f \in \mathcal{O}_M(U)$. We can write f in the form

$$f = f_0 + \sum_i f_i \xi_i + \sum_{i,j} f_{ij} \xi_i \xi_j + \dots,$$

where $f_{ij} \dots$ are some holomorphic functions on U . For $p \in U$ we will denote by $f(p)$ the value of f_0 at p .

Let $X \in T_x(M, \mathcal{O}_M)$. There is a neighborhood (U, \mathcal{O}_M) of the point x and a vector field $v_X \in \mathfrak{v}(U, \mathcal{O}_M)$ such that $\text{ev}_x(v_X) = X$. We can consider X as a linear function on $(\mathcal{O}_M)_x$. Namely, $X(f_x) := (v_X(f_x))(x)$, where $f_x \in (\mathcal{O}_M)_x$.

Let $\mu = (\mu_1, \mu_2) : (G, \mathcal{O}_G) \times (M, \mathcal{O}_M) \rightarrow (M, \mathcal{O}_M)$ be an action. Then there is a homomorphism of the Lie superalgebras $\tilde{\mu} : \mathfrak{g} \rightarrow \mathfrak{v}(M, \mathcal{O}_M)$, given by the formula $X \mapsto (\varepsilon \times \text{id})_2 \circ (X \oplus 0) \circ \mu_2$, where $X \in \mathfrak{g}$. An action μ is called *transitive* if the mapping $\text{ev}_x \circ \tilde{\mu}$ is surjective for all $x \in M$, see [4]. In this case the supermanifold (M, \mathcal{O}_M) is called (G, \mathcal{O}_G) -*homogeneous*. A supermanifold (M, \mathcal{O}_M) is called *homogeneous*, if it possesses a transitive action of some Lie supergroup.

Let $\phi = (\phi_1, \phi_2) : (M, \mathcal{O}_M) \rightarrow (M_1, \mathcal{O}_{M_1})$ be a morphism of supermanifolds. Denote by $(d\phi)_x$ the differential of ϕ at $x \in M$ (see [2, 3]). Let $\dim(M, \mathcal{O}_M) = n|m$, $\dim(M_1, \mathcal{O}_{M_1}) = k|l$. The morphism $\phi : (M, \mathcal{O}_M) \rightarrow (M_1, \mathcal{O}_{M_1})$ is called a *submersion* at $p \in M$ if $n \geq k$, $m \geq l$ and there exist two neighborhoods (U, \mathcal{O}_M) of p and (V, \mathcal{O}_{M_1}) of $q = \phi_1(p)$ with the coordinates (x_i, ξ_j) and, respectively, (y_s, η_t) such that $\phi_2|(U, \mathcal{O}_M)$ is given by the formulas:

$$\phi_2(y_i) = x_i, \quad i = 1, \dots, k, \quad \phi_2(\eta_j) = \xi_j, \quad j = 1, \dots, l. \quad (1)$$

This definition is equivalent to the requirement that the mapping $(d\phi)_p$ is surjective (see [2, 3]). A morphism $\phi = (\phi_1, \phi_2) : (M, \mathcal{O}_M) \rightarrow (M_1, \mathcal{O}_{M_1})$ is called an *immersion* at $p \in M$, if $n < k$, $m < l$, and there are neighborhoods (U, \mathcal{O}_M) and (V, \mathcal{O}_{M_1}) of p and $q = \phi_1(p)$ with coordinates (x_i, ξ_j) and, respectively, (y_s, η_t) , such that the morphism $\phi_2|(U, \mathcal{O}_M)$ is given by the formulas:

$$\begin{aligned} \phi_2(y_i) &= x_i, \quad i = 1, \dots, m, & \phi_2(y_i) &= 0, \quad i > m, \\ \phi_2(\eta_j) &= \xi_j, \quad j = 1, \dots, n, & \phi_2(\eta_j) &= 0, \quad j > n. \end{aligned} \quad (2)$$

This definition is equivalent to the requirement that the mapping $(d\phi)_p$ is injective (see [2, 3]).

Let (M, \mathcal{O}_M) be a supermanifold. A supermanifold (U, \mathcal{O}_M) , where U is an open subset in M , is called an *open subsupermanifold* of (M, \mathcal{O}_M) . Suppose that $M \subset M_1$ is a topological subspace and denote by $\phi_1 : M \rightarrow M_1$ the embedding. A supermanifold (M, \mathcal{O}_M) is called a *subsupermanifold* of a supermanifold (M_1, \mathcal{O}_{M_1}) if there is a morphism $\phi : (M, \mathcal{O}_M) \rightarrow (M_1, \mathcal{O}_{M_1})$, such that the differential $(d\phi)_p$ is injective at every point $p \in M$ and the first component of ϕ coincides with ϕ_1 . In this case we will use the notation $(M, \mathcal{O}_M) \subset (M_1, \mathcal{O}_{M_1})$.

Let $(K, \mathcal{O}_K) \subset (M, \mathcal{O}_M)$ be a subsupermanifold, $\mathcal{I} \subset \mathcal{O}_M$ the sheaf of ideals corresponding to (K, \mathcal{O}_K) , and $\mathcal{J} \subset \mathcal{O}_{G \times M}$ the sheaf of ideals corresponding to $(G \times K, \mathcal{O}_{G \times K}) \subset (G \times M, \mathcal{O}_{G \times M})$ (see [2]). We will say that (K, \mathcal{O}_K) is μ -*invariant* if the following conditions hold:

1. $\mu_1(G, K) \subset K$,
2. $\mu_2(\mathcal{I}) \subset (\mu_1)_*\mathcal{J}$.

A *Lie subsupergroup* of a Lie supergroup (G, \mathcal{O}_G) is a subsupermanifold $(H, \mathcal{O}_H) \subset (G, \mathcal{O}_G)$, such that $e \in H$ and (H, \mathcal{O}_H) is ν - and ι -invariant.

If $\phi : (M, \mathcal{O}_M) \rightarrow (M_1, \mathcal{O}_{M_1})$ is a morphism of supermanifolds and $(N, \mathcal{O}_N) \subset (M, \mathcal{O}_M)$ is a subsupermanifold, we denote by $\phi|_N$ the composition:

$$(N, \mathcal{O}_N) \hookrightarrow (M, \mathcal{O}_M) \xrightarrow{\Phi} (M_1, \mathcal{O}_{M_1}).$$

Let (G, \mathcal{O}_G) be a Lie supergroup. To each point $g \in G$ assign a morphism $\hat{g} = (\hat{g}_1, \hat{g}_2) : (\text{pt}, \mathbb{C}) \rightarrow (G, \mathcal{O}_G)$. Namely, let $\hat{g}_1(\text{pt}) = g$ and define $\hat{g}_2 : \mathcal{O}_G \rightarrow (\hat{g}_1)_*(\mathbb{C})$ by $\hat{g}_2(f_x) = 0$ for $f_x \in (\mathcal{O}_G)_x$ if $x \neq g$ and $\hat{g}_2(f_g) = f_g(g)$ for $f_g \in (\mathcal{O}_G)_g$. Denote by l_g , $g \in G$, the composition of the morphisms

$$(G, \mathcal{O}_G) \xrightarrow{\sim} (\text{pt}, \mathbb{C}) \times (G, \mathcal{O}_G) \xrightarrow{\hat{g} \times \text{id}} (G, \mathcal{O}_G) \times (G, \mathcal{O}_G) \xrightarrow{\nu} (G, \mathcal{O}_G).$$

and define r_g in a similar way. Since there exists the inverse morphism $l_{g^{-1}}$, the morphism l_g is an automorphism of the supermanifold (G, \mathcal{O}_G) , and the same is true for r_g . We refer the reader to [2, 3] for the definition of a superdomain and the proof of the following inverse function theorem.

Theorem 1. *Let (U, \mathcal{O}_U) and (V, \mathcal{O}_V) be two superdomains with coordinate systems (x_i, ξ_j) and (y_s, η_t) . Let $\phi : (U, \mathcal{O}_U) \rightarrow (V, \mathcal{O}_V)$ be a morphism and let $u \in U$. Then the following conditions are equivalent:*

- ϕ is an isomorphism in some neighborhood of u ;
- $(d\phi)_u$ is an isomorphism.

Given supermanifolds (M_i, \mathcal{O}_{M_i}) , $i = 1, \dots, n$, we will denote by $\text{pr}_{M_i}^{M_1 \times \dots \times M_n}$ the natural projection $(M_1, \mathcal{O}_{M_1}) \times \dots \times (M_n, \mathcal{O}_{M_n}) \rightarrow (M_i, \mathcal{O}_{M_i})$.

2. The structure of a supermanifold on G/H

Let $(K, \mathcal{O}_K) \subset (U, \mathcal{O}_G|_U)$ be a subsupermanifold of an open subsupermanifold $(U, \mathcal{O}_G|_U)$ in a Lie supergroup (G, \mathcal{O}_G) , \mathcal{I}_K the corresponding sheaf of ideals and φ is an isomorphism of (G, \mathcal{O}_G) . We will denote by $(\varphi(K), \mathcal{O}_{\varphi(K)})$ the subsupermanifold of $(\varphi(U), \mathcal{O}_G|_{\varphi(U)})$, where $\varphi(K) := \varphi_1(K)$, $\mathcal{O}_{\varphi(K)} := (\mathcal{O}_G|_{\varphi(U)})/(\varphi^{-1})_2(\mathcal{I}_K)$. Sometimes we will use the notation (gK, \mathcal{O}_{gK}) for $(l_g(K), \mathcal{O}_{l_g(K)})$. The subsupermanifold (Kg, \mathcal{O}_{Kg}) is defined analogously. The following proposition is well known.

† Let $\varphi = (\varphi_1, \varphi_2) : (M, \mathcal{O}_M) \rightarrow (N, \mathcal{O}_N)$ be a morphism of supermanifolds. Assume that $\varphi_1 : M \rightarrow N$ is a homeomorphism and φ is a local isomorphism. Then φ is an isomorphism.

□

Let (N, \mathcal{O}_N) and (S, \mathcal{O}_S) be two subsupermanifolds of (M, \mathcal{O}_M) . The subsupermanifold (S, \mathcal{O}_S) is called *transversal to the subsupermanifold (N, \mathcal{O}_N) at a point $x \in S \cap N$* , if $T_x(M, \mathcal{O}_M) = T_x(N, \mathcal{O}_N) \oplus T_x(S, \mathcal{O}_S)$.

Theorem 2. *Let (G, \mathcal{O}_G) be a Lie supergroup and (H, \mathcal{O}_H) be a Lie subsupergroup of (G, \mathcal{O}_G) . Suppose that $(S', \mathcal{O}_{S'})$ is a transversal subsupermanifold to (H, \mathcal{O}_H) at the point e (the identity element of G). Then there is a subsupermanifold (S, \mathcal{O}_S) , such that $\nu|_{S \times H} : (S, \mathcal{O}_S) \times (H, \mathcal{O}_H) \rightarrow (G, \mathcal{O}_G)$ is an isomorphism of $(S, \mathcal{O}_S) \times (H, \mathcal{O}_H)$ onto an open subsupermanifold $(U, \mathcal{O}_G) \subset (G, \mathcal{O}_G)$.*

Proof. Let $\dim(G, \mathcal{O}_G) = n|m$. First we will show that there exists a transversal subsupermanifold to (H, \mathcal{O}_H) at e . From the definition of a subsupermanifold it follows that there is a superdomain $(V, \mathcal{O}_V) \subset (G, \mathcal{O}_G)$ containing e , such that the subsupermanifold (H, \mathcal{O}_H) is given by the equations $x_i = 0$, $i \in \Gamma_1$, and $\xi_j = 0$, $j \in \Gamma_2$, where $\Gamma_1 \subset \{1, \dots, n\}$, $\Gamma_2 \subset \{1, \dots, m\}$. Denote by $(S', \mathcal{O}_{S'})$ the subsupermanifold in (V, \mathcal{O}_V) given by the equations $x_i = 0$, $i \notin \Gamma_1$, and $\xi_j = 0$, $j \notin \Gamma_2$. Obviously, $(S', \mathcal{O}_{S'})$ is transversal to (H, \mathcal{O}_H) .

Since $\nu|_{S \times H} \circ (\varepsilon \times \text{id}) = \text{id}$ and $\nu|_{S \times H} \circ (\text{id} \times \varepsilon) = \text{id}$, it follows that the differential $(d\nu|_{S \times H})_{(e,e)} : T_e(S, \mathcal{O}_S) \oplus T_e(H, \mathcal{O}_H) = T_e(G, \mathcal{O}_G) \rightarrow T_e(G, \mathcal{O}_G)$ is precisely the mapping

$$(v_s, v_h) \mapsto v_s + v_h. \quad (3)$$

Now it is easy to see that $(d\nu|_{S \times H})_{(e,e)}$ is an isomorphism. From Theorem 1, it follows that there are superdomains $(S, \mathcal{O}_S) \subset (S', \mathcal{O}_{S'})$, $(H', \mathcal{O}_{H'}) \subset (H, \mathcal{O}_H)$ and $(W, \mathcal{O}_G) \subset (G, \mathcal{O}_G)$, such that $\nu|_{S \times H'} : (S, \mathcal{O}_S) \times (H', \mathcal{O}_{H'}) \rightarrow (W, \mathcal{O}_G)$ is an isomorphism.

Fix a point $h \in H$. By the associativity axiom for Lie supergroups we have the following commutative diagram:

$$\begin{array}{ccc} (S, \mathcal{O}_S) \times (H', \mathcal{O}_{H'}) & \xrightarrow{id \times r_h|_H} & (S, \mathcal{O}_S) \times (H'h, \mathcal{O}_H) \\ \nu|_{S \times H'} \downarrow & & \downarrow \nu|_{S \times H'h} \\ (G, \mathcal{O}_G) & \xrightarrow{r_h} & (G, \mathcal{O}_G) \end{array}.$$

In other words, the morphism ν , restricted to $(S, \mathcal{O}_S) \times (H'h, \mathcal{O}_H)$, is equal to $r_h \circ \nu|_{S \times H'} \circ (id \times r_h|_H)^{-1}$. We have

$$(S, \mathcal{O}_S) \times (H, \mathcal{O}_H) = \bigcup_{h \in H} \{(S, \mathcal{O}_S) \times (H'h, \mathcal{O}_H)\}.$$

Therefore $\nu|_{S \times H}$ is a local isomorphism. By a well known argument from the geometric theory of homogeneous spaces we may assume that $(\nu|_{S \times H})_1 : S \times H \rightarrow G$ is a homeomorphism. Using Lemma 1, we get that $\nu|_{S \times H} : (S, \mathcal{O}_S) \times (H, \mathcal{O}_H) \rightarrow (U, \mathcal{O}_U)$ is an isomorphism. \square

Now we will give the definition of a supermanifold with the underlying manifold G/H , corresponding to a Lie supergroup (G, \mathcal{O}_G) and a subsupergroup $(H, \mathcal{O}_H) \subset (G, \mathcal{O}_G)$. Let $p_1 : G \rightarrow G/H$, $g \mapsto gH$ be the natural mapping. Fix a transversal subsupermanifold $(S, \mathcal{O}_S) \subset (G, \mathcal{O}_G)$ to (H, \mathcal{O}_H) at the point e so that $\nu|_{S \times H}$ is an isomorphism (see Theorem 2). The mapping p_1 maps S homeomorphically onto a domain $V \subset G/H$. Denote by \mathcal{O}_V the sheaf $(p_1)_*(\mathcal{O}_S)$ and identify (S, \mathcal{O}_S) with (V, \mathcal{O}_V) . Recall that the open subsupermanifold

$(U, \mathcal{O}_G) \subset (G, \mathcal{O}_G)$ is defined in Theorem 2. Let $p_U = (p_1, (p_U)_2)$ be such a morphism that the following diagram is commutative:

$$\begin{array}{ccc} (S, \mathcal{O}_S) \times (H, \mathcal{O}_H) & \xrightarrow{\nu|_{S \times H}} & (U, \mathcal{O}_G) \\ \text{pr}_V^{S \times H} \downarrow & & \downarrow p_U \\ (V, \mathcal{O}_V) & \xlongequal{\quad} & (V, \mathcal{O}_V) \end{array} \quad .$$

We can define (gV, \mathcal{O}_{gV}) analogously. Namely, it is clear that $\nu|_{gS \times H} : (gS, \mathcal{O}_{gS}) \times (H, \mathcal{O}_H) \rightarrow (gU, \mathcal{O}_{gU})$ is an isomorphism and $p_1 : gS \rightarrow gV$ is a homeomorphism. Denote by \mathcal{O}_{gV} the sheaf $(p_1)_*(\mathcal{O}_{gS})$ on gV and identify the supersubmanifold (gS, \mathcal{O}_{gS}) with (gV, \mathcal{O}_{gV}) . Define a morphism $p_{gU} : (gU, \mathcal{O}_G) \rightarrow (gV, \mathcal{O}_{gV})$ by

$$p_{gU} = \text{pr}_{gV}^{gS \times H} \circ (\nu|_{gS \times H})^{-1}.$$

Now we need the following lemma.

‡ Let (W, \mathcal{O}_W) be a superdomain, (F, \mathcal{O}_F) a supermanifold, and $(M, \mathcal{O}_M) = (W, \mathcal{O}_W) \times (F, \mathcal{O}_F)$. Denote by (x_i) all, i.e. even and odd, coordinates on (W, \mathcal{O}_W) . Then any function $f \in \mathcal{O}_M(M)$ can be written in the form $f = \sum_j h_j g_j$, where (h_j) is a maximal independent system of polynomials in x_i and $g_j \in \mathcal{O}_F(F)$.

Proof. Fix a coordinate neighborhood (N, \mathcal{O}_F) of the supermanifold (F, \mathcal{O}_F) . The function $f|_{W \times N}$ has the form $\sum_j h_j t_j^N$, where $t_j^N \in \mathcal{O}_F(N)$. Let $(\tilde{N}, \mathcal{O}_F)$ be another coordinate neighborhood of (F, \mathcal{O}_F) , such that $N \cap \tilde{N} \neq \emptyset$. Then the function $f|_{W \times \tilde{N}}$ has the form

$$f = \sum_j h_j t_j^{\tilde{N}},$$

where $t_j^{\tilde{N}} \in \mathcal{O}_F(\tilde{N})$. It is obvious that $t_j^N = \tilde{t}_j^{\tilde{N}}$ in $N \cap \tilde{N}$. Now we choose an atlas $\{(N, \mathcal{O}_F)\}$ on the supermanifold (F, \mathcal{O}_F) . The functions t_j^N are holomorphic in all coordinate neighborhoods of the chosen atlas and coincide on their intersections. It follows that there are $g_j \in H^0(F, \mathcal{O}_F)$ such that $g_j|_N = t_j^N$, and so we can write $f = \sum_j h_j g_j$. \square

Let $W \subset G/H$ be an open set. A function $f \in \mathcal{O}_G(p_1^{-1}(W))$ is called (H, \mathcal{O}_H) -right invariant if $(\nu|_{G \times H})_2(f) = (\text{pr}_{p_1^{-1}(W)}^{p_1^{-1}(W) \times H})_2(f)$ (see [1]).

‡ Let W be an open set in gV , $f \in \mathcal{O}_G(p_{gU}^{-1}(W))$. Then $f \in \text{Im}(p_{gU})_2$ if and only if f is a (H, \mathcal{O}_H) -right invariant function.

Proof. By construction, we have $\text{Im}(p_{gU})_2 = ((\nu|_{gS \times H})^{-1})_2((\text{pr}_{gS}^{gS \times H})_2(\mathcal{O}_{gS}))$. By associativity of the multiplication in (G, \mathcal{O}_G) , the following diagram is commutative:

$$\begin{array}{ccc} (gU, \mathcal{O}_G) \times (H, \mathcal{O}_H) & \xrightarrow{\nu|_{G \times H}} & (gU, \mathcal{O}_G) \\ \nu|_{gS \times H} \times \text{id} \uparrow & & \uparrow \nu|_{gS \times H} \\ (gS, \mathcal{O}_{gS}) \times (H, \mathcal{O}_H) \times (H, \mathcal{O}_H) & \xrightarrow{\text{id} \times \nu|_{H \times H}} & (gS, \mathcal{O}_{gS}) \times (H, \mathcal{O}_H) \end{array} \quad (4)$$

Now we can see that if $f \in \text{Im}(p_{gU})_2$ then f is a (H, \mathcal{O}_H) -right invariant function. Indeed, in this case $(\nu|_{gS \times H})_2(f) \in (\text{pr}_{gS}^{gS \times H})_2(\mathcal{O}_{gS})$ and

$$(\text{id}_2 \times (\nu|_{H \times H})_2)((\nu|_{gS \times H})_2(f)) = (\text{pr}_{gS}^{gS \times H \times H})_2((\nu|_{gS \times H})_2(f)).$$

Applying the isomorphism $(\nu|_{gS \times H} \times \text{id})^{-1}$ to the right hand side, we get:

$$(\nu|_{gS \times H} \times \text{id})_2^{-1} \circ (\text{pr}_{gS}^{gS \times H \times H})_2((\nu|_{gS \times H})_2(f)) = (\text{pr}_{gU}^{gU \times H})_2(f).$$

By the commutativity of (4), we obtain:

$$(\nu|_{G \times H})_2(f) = (\text{pr}_{p_1^{-1}(W)}^{p_1^{-1}(W) \times H})_2(f).$$

Conversely, let $f \in \mathcal{O}_G(p_{gU}^{-1}(W))$ be a (H, \mathcal{O}_H) -right invariant function. Without loss of generality assume that (gS, \mathcal{O}_{gS}) is a coordinate neighborhood. In the present setting, there will be no confusion to denote the even and odd coordinates by the same letters (x_i) . By Lemma 2, we have

$$(\nu|_{gS \times H})_2(f) = \sum h_i g_i,$$

where (h_i) is a maximal independent system of monomials in x_i , $g_i \in \mathcal{O}_H(H)$. By the definition of a (H, \mathcal{O}_H) -right invariant function and the commutativity of (4), we get

$$(\text{id} \times \nu|_{H \times H})_2(\sum h_i g_i) = (\text{pr}_{gS \times H}^{gS \times H \times H})_2(\sum h_i g_i) = \sum (\text{pr}_{gS}^{gS \times H \times H})_2(h_i) (\text{pr}_{H_1}^{gS \times H \times H})_2(g_i),$$

where $\text{pr}_{H_1}^{gS \times H \times H}$ is the projection onto the second factor. On the other hand,

$$(\text{id} \times \nu|_{H \times H})_2(\sum h_i g_i) = \sum (\text{pr}_{gS}^{gS \times H \times H})_2(h_i) (\text{pr}_{H \times H}^{gS \times H \times H})_2((\nu|_{H \times H})_2(g_i)).$$

The independence of (h_i) implies that $(\text{pr}_{H_1}^{H \times H})_2(g_i) = (\nu|_{H \times H})_2(g_i)$, where $\text{pr}_{H_1}^{H \times H}$ is the projection onto the first factor. Equivalently, the functions g_i are (H, \mathcal{O}_H) -right invariant. We have reduced our assertion to the following one.

(*) *If a function $g \in \mathcal{O}_H(H)$ is (H, \mathcal{O}_H) -right invariant, then $g = \text{const}$.*

Proof of ().* As above, let $\text{pr}_{H_1}^{H \times H} : (H, \mathcal{O}_H) \times (H, \mathcal{O}_H) \rightarrow (H, \mathcal{O}_H)$ denote the projection onto the first factor. By the definition of a (H, \mathcal{O}_H) -right invariant function, we have $(\text{pr}_{H_1}^{H \times H})_2(g) = (\nu|_{H \times H})_2(g)$. Now we get

$$g(e) = (\varepsilon \times \text{id})_2((\text{pr}_{H_1}^{H \times H})_2(g)) = (\varepsilon \times \text{id})_2((\nu|_{H \times H})_2(g)) = g,$$

where the last equality follows from the identity axiom $\nu \circ (\varepsilon \times \text{id}) = \text{id}$. Therefore $g = g(e)$ showing (*). This completes the proof of Lemma 2. \square

Theorem 3. *The charts (gV, \mathcal{O}_{gV}) constitute a holomorphic atlas on G/H .*

Proof. Suppose that $g_1V \cap g_2V \neq \emptyset$. Let us prove that there is a morphism $\Psi_{g_1V, g_2V} : (g_1V \cap g_2V, \mathcal{O}_{g_1V}) \rightarrow (g_1V \cap g_2V, \mathcal{O}_{g_2V})$ such that $p_{g_2U} = \Psi_{g_1V, g_2V} \circ p_{g_1U}$. Obviously, $(\Psi_{g_1V, g_2V})_1 = \text{id}$. Let us define the second component $(\Psi_{g_1V, g_2V})_2$.

If $f \in \mathcal{O}_{g_2V|_{g_1V \cap g_2V}}$ then $(p_{g_2U})_2(f) \in \mathcal{O}_G|_{g_1U \cap g_2U}$. By Lemma 2, the function $(p_{g_2U})_2(f)$ is (H, \mathcal{O}_H) -right invariant. Therefore $(p_{g_2U})_2(f) \in \text{Im}((p_{g_1U})_2)$. By construction, the map

$(p_{g_1U})_2$ is injective, and so there is a unique function $g \in \mathcal{O}_{g_1V|g_1V \cap g_2V}$, such that $(p_{g_1U})_2(g) = (p_{g_2U})_2(f)$. We put $(\Psi_{g_1V, g_2V})_2(f) := g$. The cocycle condition is obviously fulfilled. \square

We denote by $(G/H, \mathcal{O}_{G/H})$ the supermanifold defined by the holomorphic atlas constructed above. From the definition of transition functions between the charts (gV, \mathcal{O}_{gV}) , we get a morphism $p : (G, \mathcal{O}_G) \rightarrow (G/H, \mathcal{O}_{G/H})$ with $p|_{gU} = p_{gU}$ for all $g \in G$. Now we can generalize Lemma 2.

† Let $W \subset G/H$ be an open set and $f \in \mathcal{O}_G(p_1^{-1}(W))$. Then $f \in \text{Im}(p_2)$ if and only if f is (H, \mathcal{O}_H) -right invariant. \square

We now define an action of the Lie supergroup (G, \mathcal{O}_G) on the supermanifold $(G/H, \mathcal{O}_{G/H})$, which will be used in the proof of the main result of this section (Theorem 4). Denote by $i_g : (gV, \mathcal{O}_{gV}) = (gS, \mathcal{O}_{gS}) \rightarrow (gU, \mathcal{O}_G)$ the natural embedding. Let α_{gV} be the following composition:

$$(G, \mathcal{O}_G) \times (gV, \mathcal{O}_{gV}) \xrightarrow{\text{id} \times i_g} (G, \mathcal{O}_G) \times (gU, \mathcal{O}_G) \xrightarrow{\nu} (G, \mathcal{O}_G) \xrightarrow{p} (G/H, \mathcal{O}_{G/H}). \quad (5)$$

Suppose $g_1V \cap g_2V \neq \emptyset$. We claim that $\alpha_{g_1V}|_{G \times g_1V \cap g_2V} = \alpha_{g_2V}|_{G \times g_1V \cap g_2V}$. Obviously, $(\alpha_{g_1V})_1|_{G \times (g_1V \cap g_2V)} = (\alpha_{g_2V})_1|_{G \times (g_1V \cap g_2V)}$, and so we only have to prove our equality for the second components of the morphisms α_{g_1V} and α_{g_2V} . Let $W \subset G/H$ be an open set and $f \in \mathcal{O}_{G/H}(W)$. It suffices to check that

$$(\text{id}_2 \times (i_{g_1})_2)(\nu_2 \circ p_2(f)) = (\text{id}_2 \times (i_{g_2})_2)(\nu_2 \circ p_2(f)). \quad (6)$$

From the associativity of the multiplication in (G, \mathcal{O}_G) and from Lemma 3, it follows that

$$(\text{id}_2 \times (\nu|_{G \times H})_2)(\nu_2 \circ p_2(f)) = (\text{id}_2 \times (\text{pr}_G^{G \times H})_2)(\nu_2 \circ p_2(f)).$$

Let (O, \mathcal{O}_G) and (K, \mathcal{O}_G) be such open sets in (G, \mathcal{O}_G) that $O \times K \subset \nu_1^{-1}(p_1^{-1}(W))$. We assume that (O, \mathcal{O}_G) is a coordinate neighborhood with even and odd coordinates (y_j) and $K = p_1^{-1}(N)$, where N is an open set in G/H . Denote by (t_i) a maximal independent system of monomials in y_j . By Lemma 2 we can write the function $\nu_2 \circ p_2(f)|_{O \times K}$ in the form $\sum t_i s_i$, where $s_i \in \mathcal{O}_G(K)$. We have in $O \times K \times H$

$$\begin{aligned} \sum t_i (\text{pr}_G^{G \times H})_2(s_i) &= (\text{id}_2 \times (\text{pr}_G^{G \times H})_2)(\sum t_i s_i) = (\text{id}_2 \times (\text{pr}_G^{G \times H})_2)(\nu_2 \circ p_2(f)) = \\ &= (\text{id}_2 \times (\nu|_{G \times H})_2)(\nu_2 \circ p_2(f)) = (\text{id}_2 \times (\nu|_{G \times H})_2)(\sum t_i s_i) = \sum t_i (\nu|_{G \times H})_2(s_i). \end{aligned}$$

Now the equality $(\text{pr}_G^{G \times H})_2(s_i) = (\nu|_{G \times H})_2(s_i)$ follows from the independence of the system (t_i) . In other words, we get that the functions s_i are (H, \mathcal{O}_H) -right invariant. By Lemma 3 there are functions $h_i \in \mathcal{O}_{G/H}(N)$, such that $p_2(h_i) = s_i$. Moreover,

$$(\text{id}_2 \times (i_{g_1})_2)(\nu_2 \circ p_2(f)|_{O \times K}) = (\text{id}_2 \times (i_{g_1})_2)(\sum t_i s_i) = \sum t_i (i_{g_1})_2(s_i) = \sum t_i (i_{g_1})_2(p_2(h_i)).$$

Now we use the trivial equality $p \circ i_g = \text{id}$ for all $g \in G$.

$$\sum t_i (i_{g_1})_2(p_2(h_i)) = \sum t_i h_i.$$

Similarly, we obtain

$$(\text{id}_2 \times (i_{g_2})_2)(\nu_2 \circ p_2(f)|_{O \times K}) = \sum t_i h_i.$$

So we have shown that

$$(\text{id}_2 \times (i_{g_1})_2)(\nu_2 \circ p_2(f))|_{O \times N} = (\text{id}_2 \times (i_{g_2})_2)(\nu_2 \circ p_2(f))|_{O \times N}.$$

This implies (6), and so we get a morphism α , such that $\alpha|_{G \times gV} = \alpha_{gV}$. Clearly, α is an action on the supermanifold G/H . We have proved the following theorem.

Theorem 4. *There exists a supermanifold $(G/H, \mathcal{O}_{G/H})$, such that the natural action of G on G/H induces a transitive action of (G, \mathcal{O}_G) on $(G/H, \mathcal{O}_{G/H})$. The action of the Lie supergroup (G, \mathcal{O}_G) on $(G/H, \mathcal{O}_{G/H})$ is given by (5). \square*

3. Stationary Lie subsupergroup

Let $\mu : (G, \mathcal{O}_G) \times (M, \mathcal{O}_M) \rightarrow (M, \mathcal{O}_M)$ be an action of a Lie supergroup (G, \mathcal{O}_G) on a supermanifold (M, \mathcal{O}_M) and let \mathfrak{g} be the Lie superalgebra of the Lie supergroup (G, \mathcal{O}_G) . Denote by $\mu_x : (G, \mathcal{O}_G) \rightarrow (M, \mathcal{O}_M)$, $x \in M$, the composition of morphisms

$$(G, \mathcal{O}_G) \times (\text{pt}, \mathbb{C}) \xrightarrow{\text{id} \times \hat{x}} (G, \mathcal{O}_G) \times (M, \mathcal{O}_M) \xrightarrow{\mu} (M, \mathcal{O}_M),$$

where $\hat{x} = (\hat{x}_1, \hat{x}_2) : (\text{pt}, \mathbb{C}) \rightarrow (M, \mathcal{O}_M)$, $\hat{x}_1(\text{pt}) = x$, $\hat{x}_2(f) = f(x)$, $f \in \mathcal{O}_M$. Also, let $\bar{l}_g : (M, \mathcal{O}_M) \rightarrow (M, \mathcal{O}_M)$, $g \in G$, be the composition of morphisms

$$(M, \mathcal{O}_M) \simeq (\text{pt}, \mathbb{C}) \times (M, \mathcal{O}_M) \xrightarrow{\hat{g} \times \text{id}} (G, \mathcal{O}_G) \times (M, \mathcal{O}_M) \xrightarrow{\mu} (M, \mathcal{O}_M),$$

where \hat{g} was defined in Section 1.

† We have $\text{ev}_x \circ \tilde{\mu}(X) = (\text{d}\mu_x)_e(\text{ev}_e(X))$, $X \in \mathfrak{g}$. The action μ is transitive if and only if μ_x is a submersion at $e \in G$ for all $x \in M$.

Proof. The second assertion follows from the first one. Let us prove the first assertion. By definition we get

$$\text{ev}_x(\tilde{\mu}(X))(f) = (\tilde{\mu}(X)(f))(x), \quad (\text{d}\mu_x)_e(\text{ev}_e(X))(f) = (X((\mu_x)_2(f)))(e)$$

for all $f \in (\mathcal{O}_M)_x$. Therefore,

$$\begin{aligned} \text{ev}_x(\tilde{\mu}(X))(f) &= (\text{id}_2 \times \hat{x}_2)((\varepsilon_2 \times \text{id}_2)((X \oplus 0) \circ \mu_2(f))) = \\ &= (\varepsilon_2 \times \text{id}_2)((\text{id}_2 \times \hat{x}_2)((X \oplus 0) \circ \mu_2(f))) = \\ &= \varepsilon_2(X((\text{id}_2 \times \hat{x}_2) \circ \mu_2(f))) = \varepsilon_2(X((\mu_x)_2(f))) = (\text{d}\mu_x)_e(X_e)(f). \square \end{aligned}$$

The following lemma is a consequence of the axioms of action.

† We have $\mu_x \circ r_g = \mu_{gx}$, $\mu_x \circ l_g = \bar{l}_g \circ \mu_x$. \square

Consider a superdomain $(V, \mathcal{O}_M) \subset (M, \mathcal{O}_M)$, such that x is contained in V , with even coordinates (y_i) and odd coordinates (η_j) . Suppose that x has the coordinates $y_i = 0$, $\eta_j = 0$.

Denote by (U, \mathcal{O}_G) a superdomain in $((\mu_x)_1^{-1}(V), \mathcal{O}_G)$ with even coordinates (x_s) and odd coordinates (ξ_t) . Let the morphism $\mu_x|_U$ be given by the equations:

$$(\mu_x)_2(y_i) = \phi_U^i(x_s, \xi_t), \quad (\mu_x)_2(\eta_j) = \psi_U^j(x_s, \xi_t).$$

Denote by \mathcal{I}_U the sheaf of ideals in the structure sheaf of U defined as follows: if $U \cap (\mu_x)_1^{-1}(V) \neq \emptyset$ then \mathcal{I}_U is generated by the functions $\phi_U^i(x_s, \xi_t)$, $\psi_U^j(x_s, \xi_t)$, otherwise $\mathcal{I}_U := \mathcal{O}_G|_U$. The sheafs \mathcal{I}_{U_1} and \mathcal{I}_{U_2} coincide on the intersection $U_1 \cap U_2$. Therefore there is a sheaf of ideals \mathcal{I} , such that $\mathcal{I}|_U = \mathcal{I}_U$. As usual, denote by G_x the stationary subgroup of the action μ_1 at x and consider the ringed space (G_x, \mathcal{O}_{G_x}) , where $\mathcal{O}_{G_x} = (\mathcal{O}_G/\mathcal{I})|_{G_x}$.

If μ is transitive then (G_x, \mathcal{O}_{G_x}) is a subsupermanifold of (G, \mathcal{O}_G) . Indeed, by Lemma 4 the morphism μ_x is a submersion at e . From Lemma 4 it follows that μ_x is a submersion at every point $g \in G$. By the definition of a submersion, there is a coordinate neighborhood (W, \mathcal{O}_G) of $g \in G_x$ with coordinates $(x_i; \xi_j)$, $i = 1, \dots, p$, $j = 1, \dots, q$ and a coordinate neighborhood of x with coordinates $(y_a; \eta_b)$, $a = 1, \dots, k$, $b = 1, \dots, l$, such that $\mu_x|_W$ is given by the following formulas:

$$(\mu_x)_2(y_a) = x_a, \quad a = 1, \dots, k, \quad (\mu_x)_2(\eta_b) = \xi_b, \quad b = 1, \dots, l. \quad (7)$$

Without loss of generality assume that the point x is given by the system of equations $y_a = 0$, $a = 1, \dots, k$, $\eta_b = 0$, $b = 1, \dots, l$. Then $(G_x \cap W, \mathcal{O}_{G_x})$ is isomorphic to a superdomain with coordinates x_i , $i = k+1, \dots, p$, ξ_j , $j = l+1, \dots, q$.

It is not hard to prove that $\iota^2 = \text{id}$. Indeed,

$$\nu \circ (\text{id} \times \nu) \circ (\iota^2, \iota, \text{id}) = \nu \circ (\iota^2, \nu \circ (\iota, \text{id})) = \nu \circ (\iota^2, \varepsilon) = \iota^2$$

and

$$\nu \circ (\text{id} \times \nu) \circ (\iota^2, \iota, \text{id}) = \nu \circ (\nu \times \text{id}) \circ ((\iota, \text{id}) \times \text{id}) \circ (\iota, \text{id}) = \nu \circ (\varepsilon \times \text{id}) \circ (\iota, \text{id}) = \text{id}.$$

We will use this conclusion in the proof of the following theorem.

Theorem 5. *Suppose $(G_x, \mathcal{O}_{G_x}) \subset (G, \mathcal{O}_G)$ is a subsupermanifold. Then (G_x, \mathcal{O}_{G_x}) is a Lie subsupergroup of (G, \mathcal{O}_G) .*

Proof. We must show that (G_x, \mathcal{O}_{G_x}) is ν -invariant and ι -invariant. We check first that

$$(G_x, \mathcal{O}_{G_x}) \text{ is } \nu\text{-invariant.} \quad (8)$$

Obviously, $\nu_1(G_x, G_x) = G_x$. Denote by \mathcal{J} the sheaf of ideals corresponding to the subsupermanifold $(G_x, \mathcal{O}_{G_x}) \times (G_x, \mathcal{O}_{G_x})$ of the supermanifold $(G, \mathcal{O}_G) \times (G, \mathcal{O}_G)$. We have to prove that $\nu_2(\mathcal{I}) \subset \mathcal{J}$. The functions $\phi_U^i(x_s, \xi_t)$, $\psi_U^j(x_s, \xi_t)$ generate the ideal sheaf $\mathcal{I}|_U$ and $\mathcal{I}_y \neq (\mathcal{O}_G)_y$ only for $y \in G_x \subset W := (\mu_x)_1^{-1}(V)$. Therefore it is sufficient to prove that $\nu_2(\phi_U^i(x_s, \xi_t))|_{W \times W} \subset \mathcal{J}|_{W \times W}$ and $\nu_2(\psi_U^j(x_s, \xi_t))|_{W \times W} \subset \mathcal{J}|_{W \times W}$. By the definition of the functions $\phi_U^i(x_s, \xi_t)$ and $\psi_U^j(x_s, \xi_t)$, we get $\nu_2(\phi_U^i(x_s, \xi_t)) = \nu_2((\mu_x)_2(y_i))$, $\nu_2(\psi_U^j(x_s, \xi_t)) = \nu_2((\mu_x)_2(\eta_j))$. From the axioms of action it follows that $\mu_x \circ \nu = \mu \circ (\text{id} \times \mu_x)$. Thus it suffices to prove that $(\text{id}_2 \times (\mu_x)_2) \circ \mu_2(y_i)|_{W \times W} \subset \mathcal{J}|_{W \times W}$ and $(\text{id}_2 \times (\mu_x)_2) \circ \mu_2(\eta_j)|_{W \times W} \subset \mathcal{J}|_{W \times W}$.

Denote by $\text{pr}_i : (G, \mathcal{O}_G) \times (G, \mathcal{O}_G) \rightarrow (G, \mathcal{O}_G)$ the projection on the i -th factor and by $\widetilde{(\text{pr}_i)_2(\mathcal{I})}$ the sheaf of ideals generated by $(\text{pr}_i)_2(\mathcal{I})$, $i = 1, 2$. We have $\mathcal{J} = \widetilde{(\text{pr}_1)_2(\mathcal{I})} +$

$\widetilde{(\text{pr}_2)_2(\mathcal{I})}$. Further, $[(\text{id}_2 \times (\mu_x)_2) \circ \mu_2(y_i)]|_{W \times W} = (\text{id}_2 \times (\mu_x)_2)[\mu_2(y_i)|_{W \times V}]$. Using the definition of \mathcal{I} , we get:

$$(\text{id}_2 \times (\mu_x)_2)[\mu_2(y_i)|_{W \times V}] + \widetilde{(\text{pr}_2)_2(\mathcal{I})}|_{W \times W} = (\text{pr}_1)_2((\mu_x)_2(y_i))|_{W \times W} + \widetilde{(\text{pr}_2)_2(\mathcal{I})}|_{W \times W}.$$

Now, from $(\text{pr}_1)_2((\mu_x)_2(y_i))|_{W \times W} \in \widetilde{(\text{pr}_1)_2(\mathcal{I})}|_{W \times W}$ it follows that

$$(\text{id}_2 \times (\mu_x)_2) \circ \mu_2(y_i)|_{W \times W} \in \widetilde{(\text{pr}_1)_2(\mathcal{I})}|_{W \times W} + \widetilde{(\text{pr}_2)_2(\mathcal{I})}|_{W \times W} = \mathcal{I}|_{W \times W}$$

and, by the same argument, $(\text{id}_2 \times (\mu_x)_2) \circ \mu_2(\eta_j)|_{W \times W} \subset \mathcal{I}|_{W \times W}$. This completes the proof of (8).

It remains to check that

$$(G_x, \mathcal{O}_{G_x}) \text{ is } \iota\text{-invariant.}$$

Since the inclusion $\iota_1(G_x) \subset G_x$ is obvious, we must prove that $\iota_2(\mathcal{I}) \subset \mathcal{I}$ or, in terms of generators, $\iota_2(\phi_U^i(x_s, \xi_t)) \in \mathcal{I}$ and $\iota_2(\psi_U^i(x_s, \xi_t)) \in \mathcal{I}$.

By definition of the supermanifold (G_x, \mathcal{O}_{G_x}) , the following diagram is commutative:

$$\begin{array}{ccc} (G_x, \mathcal{O}_{G_x}) & \xrightarrow{\text{pr}_x} & (x, \mathbb{C}) \\ \parallel & & \downarrow \hat{x} \\ (G_x, \mathcal{O}_{G_x}) & \xrightarrow{\mu_x} & (M, \mathcal{O}_M) \end{array} .$$

We will rather use the commutativity of the next diagram:

$$\begin{array}{ccc} (\iota_1(G_x), \mathcal{O}_{\iota_1(G_x)}) & \xrightarrow{\text{pr}_x} & (x, \mathbb{C}) \\ \parallel & & \downarrow \hat{x} \\ (\iota_1(G_x), \mathcal{O}_{\iota_1(G_x)}) & \xrightarrow{\mu_x} & (M, \mathcal{O}_M) \end{array} . \quad (9)$$

To show that (9) is commutative, note that, by the inverse element axiom of a Lie supergroup and by the axioms of an action, $\mu \circ (\iota, \mu_x) = \hat{x} \circ \text{pr}_x$ and, in particular, $\mu \circ (\iota, \mu_x)|_{G_x} = (\hat{x} \circ \text{pr}_x)|_{G_x}$. Using the equality $\mu_x|_{G_x} = \hat{x}$ and the definition of the morphism μ_x , we obtain the commutative diagram:

$$\begin{array}{ccc} (G_x, \mathcal{O}_{G_x}) & \xrightarrow{\text{pr}_x} & (x, \mathbb{C}) \\ (\iota, \hat{x}) \downarrow & & \downarrow \hat{x} \\ (\iota_1(G_x), \mathcal{O}_{\iota_1(G_x)}) \times (x, \mathbb{C}) & \xrightarrow{\mu_x} & (M, \mathcal{O}_M) \end{array} .$$

Since (ι, \hat{x}) is an isomorphism, we get the commutativity of (9).

Denote by $\mathcal{I}_{\iota(G_x)}$ the sheaf of ideals in \mathcal{O}_G corresponding to the subsupermanifold $(\iota_1(G_x), \mathcal{O}_{\iota_1(G_x)}) \subset (G, \mathcal{O}_G)$. Using (9), we have:

$$\phi_U^i(x_s, \xi_t) + \mathcal{I}_{\iota(G_x)} = \mu_x(y_i) + \mathcal{I}_{\iota(G_x)} = y_i(x) + \mathcal{I}_{\iota(G_x)} = 0 + \mathcal{I}_{\iota(G_x)} = \mathcal{I}_{\iota(G_x)}.$$

Therefore $\phi_U^i(x_s, \xi_t) \in \mathcal{I}_{\iota(G_x)}$. Similarly, $\psi_U^i(x_s, \xi_t) \in \mathcal{I}_{\iota(G_x)}$. It follows that $\mathcal{I} \subset \mathcal{I}_{\iota(G_x)} = \iota_2(\mathcal{I})$, and the equality $\iota^2 = \text{id}$ implies that $\mathcal{I} = \mathcal{I}_{\iota(G_x)}$. \square

We have seen that for a transitive action of (G, \mathcal{O}_G) on (M, \mathcal{O}_M) the ringed space (G_x, \mathcal{O}_{G_x}) is a supermanifold, which is in fact a Lie subsupergroup of (G, \mathcal{O}_G) . This Lie subsupergroup will be called *the stationary Lie subsupergroup of x* . In the last section we will prove that any homogeneous supermanifold is isomorphic to a coset space of a Lie supergroup with the structure of a supermanifold introduced above.

4. The structure of a homogeneous space

Our goal here is the following theorem.

Theorem 6. *Let (M, \mathcal{O}_M) be a (G, \mathcal{O}_G) -homogeneous supermanifold and let (H, \mathcal{O}_H) be a stationary Lie subsupergroup of a point $x \in M$. Then there is an isomorphism*

$$\beta : (G/H, \mathcal{O}_{G/H}) \rightarrow (M, \mathcal{O}_M),$$

which is (G, \mathcal{O}_G) -equivariant in the sense that the following diagram is commutative:

$$\begin{array}{ccc} (G, \mathcal{O}_G) \times (G/H, \mathcal{O}_{G/H}) & \xrightarrow{\text{id} \times \beta} & (G, \mathcal{O}_G) \times (M, \mathcal{O}_M) \\ \alpha \downarrow & & \downarrow \mu \\ (G/H, \mathcal{O}_{G/H}) & \xrightarrow{\beta} & (M, \mathcal{O}_M) \end{array} . \quad (10)$$

Proof. Let $\beta_1 : G/H \rightarrow M$, $gH \mapsto gx$, be the natural homeomorphism. Then the diagram

$$\begin{array}{ccc} (G, \mathcal{O}_G) & \xlongequal{\quad} & (G, \mathcal{O}_G) \\ p=(p_1, p_2) \downarrow & & \downarrow \mu_x=((\mu_x)_1, (\mu_x)_2), \\ (G/H, \mathcal{O}_{G/H}) & \xrightarrow{\beta_1} & (M, \mathcal{O}_M) \end{array} \quad (11)$$

is commutative. We will now construct an isomorphism of sheaves $\beta_2 : \mathcal{O}_M \rightarrow (\beta_1)_* \mathcal{O}_{G/H}$, such that $\beta = (\beta_1, \beta_2)$ is the required isomorphism.

As in Theorem 2, consider the product $(S, \mathcal{O}_S) \times (H, \mathcal{O}_H)$. By the axioms of action, the composition of morphisms

$$(S, \mathcal{O}_S) \times (H, \mathcal{O}_H) \xrightarrow{\nu|_{S \times H}} (U, \mathcal{O}_G) \xrightarrow{\mu_x} (M, \mathcal{O}_M).$$

can be written as $\mu_x \circ \nu|_{S \times H} = \mu \circ (\text{id} \times \mu_x|_H) = \mu \circ (\text{id} \times \hat{x})$. The last equality follows from the fact that $\mu_x|_H = \hat{x}$. As a result, we get

$$\mu_x \circ \nu|_{S \times H} = \mu_x \circ \text{pr}_S^{S \times H}. \quad (12)$$

The differential $(d\mu_x)_e$ is surjective, because the action μ is transitive. The differential $(d\nu|_{S \times H})_{(e, e)}$ is nondegenerate, see Theorem 2. It is easy to see that $\dim(S, \mathcal{O}_S) = \dim(M, \mathcal{O}_M)$, and so we get that $(d(\mu_x)|_S)_{(e)}$ is nondegenerate. By Theorem 1, we can assume that $\mu|_S$ is an isomorphism of (S, \mathcal{O}_S) onto some superdomain $(V, \mathcal{O}_M) \subset (M, \mathcal{O}_M)$. By (12) we

get $(\mu_x)_2(\mathcal{O}_M|_V) = ((\nu|_{S \times H})^{-1})_2(\mathcal{O}_S)$. On the other hand, by the definition of p we have $p_2(\mathcal{O}_{G/H}|_{p_1(S \times H)}) = ((\nu|_{S \times H})^{-1})_2(\mathcal{O}_S)$ and therefore $p_2(\mathcal{O}_{G/H}|_{p_1(S \times H)}) = (\mu_x)_2(\mathcal{O}_M|_V)$.

Let $\tilde{l}_g = \alpha \circ (\hat{g} \times \text{id})$. The following diagram is commutative by Lemma 4:

$$\begin{array}{ccc}
(p_1(S \times H), \mathcal{O}_{G/H}) & \xrightarrow{\tilde{l}_g} & (p_1(gS \times H), \mathcal{O}_{G/H}) \\
\uparrow p & & \uparrow p \\
(U, \mathcal{O}_G) & \xrightarrow{l_g} & (gU, \mathcal{O}_G) \\
\downarrow \mu_x & & \downarrow \mu_x \\
(V, \mathcal{O}_M) & \xrightarrow{\tilde{l}_g} & (gV, \mathcal{O}_M)
\end{array} \quad . \quad (13)$$

By the commutativity of (13), we get

$$(\mu_x)_2(\mathcal{O}_M|_{gV}) = p_2(\mathcal{O}_{G/H}|_{p_1(gU)}), \quad \text{for all } g \in G.$$

Therefore, the sheaves $(\mu_x)_2(\mathcal{O}_M)$ and $p_2(\mathcal{O}_{G/H})$ coincide locally, and it follows that $p_2(\mathcal{O}_{G/H}) = (\mu_x)_2(\mathcal{O}_M)$. Define a morphism $\beta_2 : \mathcal{O}_M \rightarrow (\beta_1)_*(\mathcal{O}_{G/H})$ by

$$\beta_2(f) = p_2^{-1} \circ (\mu_x)_2(f).$$

Obviously, the morphism $\beta = (\beta_1, \beta_2)$ is an isomorphism and the diagram (11) is commutative.

It remains to prove that diagram (10) is also commutative. By the definition of β , we get locally $\beta = (\beta_1, \beta_2) = \mu_x \circ i_g$, where i_g is defined after Lemma 3. Hence $\mu_x \circ i_g \circ p = \mu_x$ and the axioms of action yield $\mu \circ (\text{id} \times \mu_x) = \mu_x \circ \nu$. Thus we get locally:

$$\mu \circ (\text{id} \times \beta) = \mu \circ (\text{id} \times \mu_x \circ i_g) = \mu \circ (\text{id} \times \mu_x) \circ (\text{id} \times i_g) = \mu_x \circ \nu \circ (\text{id} \times i_g).$$

$$\beta \circ \alpha = \mu_x \circ i_g \circ p \circ \nu \circ (\text{id} \times i_g) = \mu_x \circ \nu \circ (\text{id} \times i_g).$$

This implies the commutativity of the diagram (10). \square

The author would like to thank her supervisor Prof. Arkady Onishchik for many helpful remarks on the paper, Prof. Dmitry Akhiezer for interesting discussions and Prof. Dr. Dr. h.c. mult. Alan T. Huckleberry for the hospitality at Ruhr-Universität Bochum.

References

- [1] *Kostant B.* Graded Manifolds, graded Lie theory, and prequantization. Lecture Notes in Mathematics 570. Berlin e.a.: Springer-Verlag, 1977. P. 177-306.
- [2] *Leites D.A.* Introduction to the theory of supermanifolds. Russian Math. Surveys 35 (1980), 1-64.
- [3] *Manin Yu. I.* Gauge field theory and complex geometry. Grundlehren der Mathematischen Wissenschaften, 289, Springer-Verlag, Berlin, 1997.
- [4] *Onishchik A.L.* Flag supermanifolds, their automorphisms und deformations. The Sophus Lie Memorial conference (Oslo, 1992), 289-302, Scand. Univ. Press, Oslo, 1994.

TVER STATE UNIVERSITY, ZHELYABOVA 33, 170 000 TVER, RUSSIA
E-mail address: VishnyakovaE@googlemail.com